

RESEARCH ARTICLE

Open Access



Long-term antihypertensive effects of far-infrared ray irradiated from wooden board in spontaneously hypertensive rats

Chien-Tsong Lin^{1,2}, Ming-Ju Lin³, Yung-Pin Chen², Ko-Chao Lee⁴, Kuo-Chin Huang⁵, Shun-Fu Chang^{6*} and Cheng-Nan Chen^{7*}

Abstract

Background: Far-infrared ray (FIR) has been widely used in promoting health and has been shown to exert beneficial effects in vascular function. The non-thermal effect of FIR has been found to play a significant role in the protective effect on some vascular-related diseases, but its protective effects and use against hypertension have not been clearly presented.

Methods: In the present study, by using a wooden board coated with FIR-irradiated materials, we evaluated the long-term antihypertensive effect on spontaneously hypertensive rats (SHRs) in the environment in contact with the FIR-irradiated wooden board. SHRs were placed on the wooden board with or without FIR radiation for 4 weeks.

Results: The systolic blood pressure (BP) of SHRs undergoing different treatments was measured weekly using a tail-cuff method. FIR radiation significantly reduced the systolic BP of the SHRs along with a decreasing plasma level of angiotensin II and an increasing plasma level of bradykinin. In addition, long-term contact of FIR did not significantly affect the BP in normotensive Wistar Kyoto rats (WKYs).

Conclusions: Our results provided the evidence based on which FIR radiation could be considered an effective non-pharmacological choice for preventing hypertension.

Keywords: Angiotensin II, Far-infrared ray, Hypertension, Spontaneously hypertensive rats

Background

Hypertension, the persistent increase of blood pressure (BP) above 140/90 mmHg, is a major risk factor for the development of cardiovascular disease, including stroke, myocardial infarction, heart and renal diseases, and cardiac mortality. Although the etiology of hypertension is complex, the renin-angiotensin system (RAS) plays a major role in the pathology of hypertension [1, 2]. Renin catalyzes the conversion of angiotensinogen to angiotensin I, which is the rate-determining step. Angiotensin-converting enzyme (ACE) then catalyzes the conversion of angiotensin I to angiotensin II, leading to the constriction of the arterial vessels and

the regulation of systemic BP [3]. Since the increased levels of angiotensin II are closely related to the development and maintenance of hypertension, the inhibition of angiotensin II formation has been widely considered a major strategy for lowering BP in patients with hypertension [4, 5]. Because sustained BP elevation may increase the incidence of atherosclerosis and stroke, it is important to develop alternative strategies for the treatment of hypertension.

The far-infrared ray (FIR) is generally defined as the electromagnetic wave in the wavelength range of 5.6–1000 μm . FIR radiation has been widely applied to a variety of fields as it has positive effects on food preservation [6], health promotion [7], and cardiovascular systems [8, 9]. Some studies indicated that FIR therapy has potentially beneficial effects in the treatment of wound healing [10], diabetes [11], tumor thermal therapy [12], chronic fatigue syndrome [13], and knee osteoarthritis [14]. In addition, several

* Correspondence: sfchang@cgmh.org.tw; cnchen@mail.nyu.edu.tw

⁶Department of Medical Research and Development, Chang Gung Memorial Hospital Chiayi Branch, No. 6, West Sec., Jiapu Rd., Puzi City, Chiayi County 613, Taiwan, ROC

⁷Department of Biochemical Science and Technology, National Chiayi University, Chiayi 600, Taiwan

Full list of author information is available at the end of the article



© 2016 Lin et al. **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

studies also indicated that FIR radiation exerts beneficial effects in the treatment of vascular diseases. FIR radiation improved impaired vascular endothelial function in patients with heart disease [15], decreased endothelial inflammation by induction of heme oxygenase-1 [16], and augmented angiogenesis in the ischemic hind limb [11]. These studies revealed that the potential effects of FIR therapy played a significant role in the protective effect on vascular function.

There is an increasing evidence that the secretion of soluble inflammatory makers play a significant role in hypertension [17, 18]. Soluble intercellular adhesion molecule 1 (sICAM-1), vascular adhesion molecule 1 (sVCAM-1), and sE-selectin are major inflammatory markers and endothelial cell adhesion molecules that regulates the binding and extravasation of leukocytes from the bloodstream to sites of inflammation [19]. The appearance of soluble inflammatory markers in the circulation is thought to be the consequence of their release from the vascular cells because of increased expression [20].

The spontaneous hypertension rats (SHRs) are the most widely used animal model in the study of hypertension-related diseases. The duration, cardiac hypertrophy, renal dysfunction, and cardiovascular complications of SHRs are similar to those observed in human hypertension. In this study, by using the wooden board coated with FIR-emitting materials, we evaluated the long-term effect on SHRs in contact with the FIR-emitting wooden board. Wistar Kyoto rats (WKYs) were used as the control group. These findings provide new insights into the treatment of hypertension using FIR radiation.

Methods

The FIR-irradiated wooden board

All of the FIR-irradiated wooden board (53 × 21.5 × 1.5 cm; length × width × thickness) samples used in this study were made by Ua Wood Floors, Inc. (Yunlin, Taiwan). The wooden boards were coated with Nano-paste (ZnO), which was manufactured by Nano-Infinity Nanotech Co., Ltd. (Taoyuan, Taiwan). Using the far-infrared ray testing system (Bruker, VERTEX 70 FT-IR) to test for far-infrared emissivity, the spectrums of the FIR-irradiated wooden board exhibited high emissivity (>0.8) between 4 ~ 14 μm wavelengths (triplicate samples were made for each wooden board, with measurements done by Industrial Technology Research Institute in Taiwan).

Animal model

The study was performed in accordance with the guidelines for laboratory animals of National Chiayi University. The protocol was approved by the Institutional Animal Care and Use Committee of National Chiayi University. All efforts were made to minimize suffering. Eight-

week-old male SHRs and WKYs were obtained from the National Laboratory Animal Breeding and Research Center of the National Science Council, Taiwan. All rats were maintained five per cage at a constant temperature (24 ± 1 °C), with a 12-h dark/light cycle and a standard rat chow diet. After the SHRs had been housed for 10 weeks, their weight ranged from 240–250 g, and systolic BP reached 180 mmHg. The rats were randomly divided into four groups ($n = 10$ per group). Group 1: SHRs on a wooden board without FIR radiation as the control group; group 2: SHRs on a wooden board with FIR radiation; group 3: SHRs on a normal wooden board with atenolol administration as the positive control group; and group 4: SHRs on an FIR-radiated wooden board with atenolol administration. In addition, the WKY rats were divided into group 1 and group 2 for comparing the SHRs in this study.

Measurement of BP

BP was measured indirectly and non-invasively every week by using the volume-oscillometric tail-cuff method (UR-5000, UEDA, Industries Co., Tokyo, Japan). To get an accurate BP, at least 7 consecutive determinations were recorded. The maximum and minimum values of BP were then eliminated, and the BP was calculated as the average of the remaining values [21].

Analysis of plasma angiotensin II, bradykinin, and ACE activity

After animal sacrifice, blood samples were collected and kept on ice. The samples were collected and centrifuged at 3000 rpm for 15 min at 4 °C. The plasma was stored at -80 °C until analysis. Both angiotensin II and bradykinin were quantified by respective ELISA kits (Angiotensin II ELISA, Cayman Chemical, Ann Arbor, MI, USA; Bradykinin ELISA, Phoenix Pharmaceuticals, Burlingame, CA, USA) as per the manufacturers' instructions. ACE activity was determined according to the colorimetric method using synthetic substrate (FAPGG) [22]. One unit (U) of ACE activity is defined as the amount of which released 1 μmol of His-Leu per minute. The specific activity of ACE is expressed as U/L.

Analysis of plasma inflammatory markers

The levels of sICAM-1, sVCAM-1, and sE-selectin in plasma were determined by using sandwich ELISA (R&D Systems, Minneapolis, MN) according to manufacturer's protocols [23].

Statistics analysis

Data were reported as the mean ± standard error of the means (SEM) and evaluated by one-way analysis of variance. Significant differences were established at $P < 0.05$.

Results

FIR-irradiated wooden board attenuates BP in SHRs

The SHRs demonstrated already established hypertension at the beginning of the study (week 0). The results from this study, as illustrated in Fig. 1a, demonstrated that at the start of the experiment, the mean values of systolic BP in all the SHRs were similar (174.5 to 179.6 mmHg). The systolic BP in the control SHRs was increased up to 193.6 mmHg at the end of the 4-week period of experiment, whereas SHRs on an FIR-irradiated wooden board started to reduce systolic BP after 1 week, and this effect continued until the 4-week experimental period with gradual reduction in systolic BP. Among the treatment, SHR administration with atenolol reduced the systolic BP more significantly at week 3 and 4 compared with the SHRs treated with FIR




Fig. 1 Changes in systolic BP of experimental rats during the 4-week period. **a** The SHRs were randomly divided into four groups ($n = 10$ per group). Group 1: SHRs on a wooden board without FIR radiation as the control group (control); group 2: SHRs on a wooden board with FIR radiation (FIR); group 3: SHRs on a normal wooden board with atenolol administration as the positive control group (atenolol); and group 4: SHRs on an FIR-irradiated wooden board with atenolol administration (FIR + atenolol). **b** The WKY rats were randomly divided into two groups ($n = 10$ per group). Group 1: WKYs on a wooden board without FIR radiation as the control group (control); and group 2: WKY rats on a wooden board with FIR radiation (FIR). Data are mean \pm SEM. * $p < 0.05$ vs. control SHRs, # $p < 0.05$ vs. SHRs with FIR irradiation

only. There were no obvious changes in the systolic BP in WKYs with or without FIR irradiation (Fig. 1b). These experiments suggested that the effect of FIR irradiation on SBP levels has specificity against hypertensive rats.

FIR irradiation inhibits plasma ACE activity in SHRs

Plasma ACE activity in the SHR untreated group was significantly higher than in the WKY group (35.4 ± 3.1 vs. 124.5 ± 8.4 U/L). FIR irradiation decreased ACE activity in the SHR untreated group to 76.7 ± 6.3 U/L in the FIR-irradiated group (Fig. 2). In addition, SHR administration with atenolol further reduced the plasma ACE activity compared with the SHRs treated with FIR only (Fig. 2).

FIR irradiation appears to inhibit angiotensin II in vivo

The effect of FIR irradiation on the systolic BP of the SHRs was associated with concomitant changes in circulating levels of angiotensin II and bradykinin. FIR irradiation decreased angiotensin II levels from 48.2 ± 5.4 pg/mL in the untreated group to 30.5 ± 3.1 pg/mL in the FIR-irradiated group (Fig. 3). In addition, SHR administration with atenolol further reduced the angiotensin II levels compared with the SHRs treated with FIR only (Fig. 3). The treatment also increased the circulating levels of bradykinin (a molecule metabolized by ACE), which increased from 1.82 ng/mL in the untreated group to 3.05 ng/mL in the FIR irradiated group (Fig. 4). Moreover, the WKY rats had the lowest angiotensin II (Fig. 3) and the highest bradykinin (Fig. 4) levels, compared to all the SHR groups. These results suggest that FIR irradiation may decrease the production of angiotensin II as






Fig. 2 Plasma ACE activity of experimental rats after the 4-week treatments. WKYs: WKYs on a wooden board without FIR radiation; CL: SHRs on a wooden board without FIR radiation; FIR: SHRs on a wooden board with FIR radiation; atenolol: SHRs on a normal wooden board with atenolol administration; FIR + atenolol: SHRs on an FIR-irradiated wooden board with atenolol administration. Data are mean \pm SEM from $n = 10$ animals per treatment group. * $p < 0.05$ vs. control SHRs, # $p < 0.05$ vs. SHRs with FIR irradiation, † $p < 0.05$ vs. WKY group



well as inhibit the degradation of bradykinin, thereby decreasing the systolic BP in SHRs.

FIR irradiation ameliorates plasma levels of inflammatory markers

We further evaluated the effect of FIR irradiation on plasma levels of sICAM-1, sVCAM-1, and sE-selectin in SHRs. SHR untreated group exhibited an increase in sICAM-1, sVCAM-1, and sE-selectin levels compared to




WKY rats (Fig. 5). However, SHRs treated with FIR showed decreased sICAM-1, sVCAM-1, and sE-selectin levels compared to untreated group (Fig. 5).

Discussion

Recent findings showed that FIR radiation might play a role in the long-term protective effect on vascular function by reducing several cardiovascular risk factors such as high BP. Our studies with FIR irradiation in SHRs demonstrated for the first time that rats placed on FIR irradiated wooden board indeed showed a systolic BP reducing effect in the hypertensive rats in a time-dependent manner. The FIR also decreased angiotensin II and increased bradykinin production in the plasma of the treated SHRs, suggesting the possible mechanisms for the systolic BP reducing effect of FIR. Thus, FIR could prove to be a novel method for management of hypertension.

SHRs are the animal model most frequently used to evaluate the in vivo antihypertensive effect of different treatments such as drugs or dietary components. Various studies performed to determine the antihypertensive effects of food-derived bioactive peptides have used SHR animals as a model system [24, 25]. To verify the validity of the experimental protocol, both a negative and a positive control group were included. The underlying pathological mechanisms of the hypertensive effect in SHRs include increased activity of the renin-angiotensin system as well as increases in vascular inflammation and oxidative stress [26, 27]. Hence, SHRs are a suitable model to study the in vivo antihypertensive efficacy of FIR irradiation. In the present study, systolic BP increased in the control SHR group during the 4-week period of the study. In addition, the systolic BP was significantly decreased in the positive control group (which received atenolol administration) when compared to the negative control group. Thus, the negative and positive control groups confirmed the validity of the animal model chosen in this study.

The increase in BP is directly due to the vasoconstrictive actions of angiotensin II produced from angiotensin I in the presence of ACE [28, 29]. Angiotensin II significantly increased the BP by the activation of angiotensin II receptors in the vascular smooth muscle cells, resulting in increased vasoconstriction, decreased renal blood flow and renal tubular sodium re-uptake, and increased vasopressin secretion [30–32]. In our present study, FIR radiation appears to act through angiotensin II, leading to lowered BP in SHRs. FIR radiation reduced the plasma and angiotensin II levels with a corresponding increase in bradykinin. The plasma levels of ACE activity tended to be positively related to BP changes, which is consistent with the earlier work that have shown that ACE levels could be used as a hypertension marker in



SHRs [33]. Our result is further confirmed by the fact that FIR-irradiated SHRs had significantly reduction of plasma ACE activity when compared to the untreated control, suggesting a role of elevated plasma ACE activity in modulating hypertension conditions. Thus, the antihypertensive effects of FIR radiation may be mediated by downregulation of ACE with a consequent reduction in the plasma levels of the vasoconstrictor angiotensin II.

It has been showed that FIR therapy has been applied to various clinical fields, particularly including vessel-related disorders [34]. FIR radiation enhanced blood flow retrieval and blood vessel formation in ischemic hind limbs by enhancing the homing of endothelial progenitor cells and lowering the oxidative stress in ischemic tissue. Direct FIR radiation could also ameliorate high-glucose-induced oxidative stress, suppress cellular senescence, and improve endothelial progenitor cell functions [11]. Although the mechanism by which FIR radiation has a potential for the treatment of vascular diseases is still unclear, the improvement of endothelial cell function is currently considered as a mechanism [35]. In addition, increasing evidence suggests FIR therapy could exert beneficial effects in the cardiovascular system through the NO-related pathway [36]. FIR therapy could upregulate eNOS expression and increase angiogenesis in an apolipoprotein E-knockout mouse model [37]. These findings suggest that FIR may provide some

novel rationales for its potential clinical impact on vascular protection.

Vascular dysfunction or damage can be evaluated by the measurement of inflammatory markers released from vascular cells, such as sICAM-1, sVCAM-1, and sE-selectin [38]. Several studies have demonstrated that the plasma levels of inflammatory markers are increased in patients with hypertension, type 2 DM, and atherosclerosis, all of which are well-established risk factors for cardiovascular diseases [39–41]. Higher levels of soluble inflammatory markers are associated with increased cardiovascular risk in hypertension, presumably because such high levels reflect chronic, low-grade vascular inflammation [18]. FIR therapy has been reported to exert a potent anti-inflammatory effect via induction of HO-1 release by human umbilical vein endothelial cells [16]. Down-regulation in these markers may confer a decrease in cardiovascular risk, although prospective data are needed to test this hypothesis. In the current study, FIR irradiation decreased levels of the inflammatory markers sICAM-1, sVCAM-1, and sE-selectin, suggesting a specific anti-inflammatory property of FIR in vivo. Since SHR untreated group has increased levels of plasma inflammatory markers which may contribute to endothelial dysfunction, it is suggested that controlling the inflammatory responses with FIR irradiation could potentially improve the vascular function and reduce inflammation.

Conclusions

This study provides evidence of the antihypertensive effect of FIR irradiation in SHR. We found that the treatment of SHR with FIR irradiation induced a decrease in BP. Since FIR substantially decreased the SHR plasma ACE activity and angiotensin II level, and increased the bradykinin level, we believe that the mechanism of the antihypertensive effect of FIR on SHR may be related to the ACE inhibitory effect. Our results suggest a possible beneficial effect of FIR on cardiovascular function and thus provide a mechanistic basis for the use of FIR as an antihypertensive remedy in folk medicine.

Abbreviations

BP: blood pressure; RAS: renin-angiotensin system; ACE: angiotensin-converting enzyme; FIR: far-infrared ray; SHR: spontaneous hypertension rat; WKY: Wistar Kyoto rat.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CTL, CNC, and SFC wrote the manuscript. CTL, CNC, and SFC designed the study and performed the statistical analysis. MJL, YPC, KCL, and KCH were involved in the acquisition of data. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by a grants from the Yunlin County Government, Taiwan (102SBIR-A07), Chang Gung Memorial Hospital Chiayi Branch (CMRPG6E0231), and by the Ministry of Science and Technology, Taiwan (NSC101-2320-B-415-003-MY3, MOST103-2320-B182A-021-MY3).

Author details

¹Center for General Education, National Formosa University, Yunlin 632, Taiwan. ²Department of Wood Based Materials and Design, National Chiayi University, Chiayi 600, Taiwan. ³Department of physical Education, Health and Recreation, National Chiayi University, Minsyong, Chiayi County 621, Taiwan. ⁴Department of Colorectal Surgery, Department of Surgery, Chang Gung Memorial Hospital; Kaohsiung Medical Center, Kaohsiung 833, Taiwan. ⁵Department of Orthopaedics, Chang Gung Memorial Hospital Chiayi Branch, Chiayi 613, Taiwan. ⁶Department of Medical Research and Development, Chang Gung Memorial Hospital Chiayi Branch, No. 6, West Sec., Jiapu Rd., Puzi City, Chiayi County 613, Taiwan, ROC. ⁷Department of Biochemical Science and Technology, National Chiayi University, Chiayi 600, Taiwan.

Received: 7 October 2014 Accepted: 5 February 2016

Published online: 08 February 2016

References

- van Vark LC, Bertrand M, Akkerhuis KM, Bruggs JJ, Fox K, Mourad JJ, et al. Angiotensin-converting enzyme inhibitors reduce mortality in hypertension: a meta-analysis of randomized clinical trials of renin-angiotensin-aldosterone system inhibitors involving 158,998 patients. *Eur Heart J*. 2012;33:2088–97.
- van Twist DJ, Kroon AA, de Leeuw PW. Angiotensin-(1-7) as a strategy in the treatment of hypertension? *Curr Opin Nephrol Hypertens*. 2014;23:480–6.
- Bernstein KE, Giani JF, Shen XZ, Gonzalez-Villalobos RA. Renal angiotensin-converting enzyme and blood pressure control. *Curr Opin Nephrol Hypertens*. 2014;23:106–12.
- Giani JF, Janjulia T, Taylor B, Bernstein EA, Shah K, Shen XZ, et al. Renal generation of angiotensin II and the pathogenesis of hypertension. *Curr Hypertens Rep*. 2014;16:477.
- Patel SK, Velkoska E, Freeman M, Wai B, Lancefield TF, Burrell LM. From gene to protein-experimental and clinical studies of ACE2 in blood pressure control and arterial hypertension. *Front Physiol*. 2014;5:227.
- Lee KT. Quality and safety aspects of meat products as affected by various physical manipulations of packaging materials. *Meat Sci*. 2010;86:138–50.
- Cho S, Shin MH, Kim YK, Seo JE, Lee YM, Park CH, et al. Effects of infrared radiation and heat on human skin aging in vivo. *J Investig Dermatol Symp Proc*. 2009;14:15–9.
- Lohr NL, Keszler A, Pratt P, Bienengraber M, Warltier DC, Hogg N. Enhancement of nitric oxide release from nitrosyl hemoglobin and nitrosyl myoglobin by red/near infrared radiation: potential role in cardioprotection. *J Mol Cell Cardiol*. 2009;47:256–63.
- Inoue S, Takemoto M, Chishaki A, Ide T, Nishizaka M, Miyazono M, et al. Leg heating using far infra-red radiation in patients with chronic heart failure acutely improves the hemodynamics, vascular endothelial function, and oxidative stress. *Intern Med*. 2012;51:2263–70.
- Toyokawa H, Matsui Y, Uhara J, Tsuchiya H, Teshima S, Nakanishi H, et al. Promotive effects of far-infrared ray on full-thickness skin wound healing in rats. *Exp Biol Med (Maywood)*. 2003;228:724–9.
- Huang PH, Chen JW, Lin CP, Chen YH, Wang CH, Leu HB, et al. Far infra-red therapy promotes ischemia-induced angiogenesis in diabetic mice and restores high glucose-suppressed endothelial progenitor cell functions. *Cardiovasc Diabetol*. 2012;11:99.
- Udagawa Y, Ishigame H, Nagasawa H. Effects of hydroxyapatite in combination with far-infrared rays on spontaneous mammary tumorigenesis in SHN mice. *Am J Chin Med*. 2002;30:495–505.
- Masuda A, Kihara T, Fukudome T, Shinsato T, Minagoe S, Tei C. The effects of repeated thermal therapy for two patients with chronic fatigue syndrome. *J Psychosom Res*. 2005;58:383–7.
- Bagnato GL, Miceli G, Atteritano M, Marino N, Bagnato GF. Far infrared emitting plaster in knee osteoarthritis: a single blinded, randomised clinical trial. *Rheumatismo*. 2012;64:388–94.
- Imamura M, Biro S, Kihara T. Repeated thermal therapy improves impaired vascular endothelial function in patients with coronary risk factors. *J Am Coll Cardiol*. 2001;38:108331088.
- Lin CC, Liu XM, Peyton K, Wang H, Yang WC, Lin SJ, et al. Far infrared therapy inhibits vascular endothelial inflammation via the induction of heme oxygenase-1. *Arterioscler Thromb Vasc Biol*. 2008;28:739–45.
- Preston RA, Ledford M, Materson BJ, Baltodano NM, Memon A, Alonso A. Effects of severe, uncontrolled hypertension on endothelial activation: soluble vascular cell adhesion molecule-1, soluble intercellular adhesion molecule-1 and von Willebrand factor. *J Hypertens*. 2002;20:871–7.
- Zhang M, Wang G, Wang A, Tong W, Zhang Y. Association of hypertension with coexistence of abnormal metabolism and inflammation and endothelial dysfunction. *Blood Press*. 2013;22:151–7.
- Sato J, Kinugasa M, Satomi-Kobayashi S, Hatakeyama K, Knox AJ, Asada Y, et al. Family with sequence similarity 5, member C (FAM5C) increases leukocyte adhesion molecules in vascular endothelial cells: implication in vascular inflammation. *PLoS One*. 2014;9:e107236.
- Tesfamariam B, DeFelicis AF. Endothelial injury in the initiation and progression of vascular disorders. *Vascul Pharmacol*. 2007;46:229–37.
- Inanaga K, Ichiki T, Matsuura H, Miyazaki R, Hashimoto T, Takeda K, et al. Resveratrol attenuates angiotensin II-induced interleukin-6 expression and perivascular fibrosis. *Hypertens Res*. 2009;32:466–71.
- García-Tejedor A, Sánchez-Rivera L, Castelló-Ruiz M, Recio I, Salom JB, Manzanares P. Novel antihypertensive lactoferrin-derived peptides produced by *Kluyveromyces marxianus*: gastrointestinal stability profile and in vivo angiotensin I-converting enzyme (ACE) inhibition. *J Agric Food Chem*. 2014;62:1609–16.
- Chen TC, Chien SJ, Kuo HC, Huang WS, Sheen JM, Lin TH, et al. High glucose-treated macrophages augment E-selectin expression in endothelial cells. *J Biol Chem*. 2011;286:25564–73.
- Xu L, Liu Y. Administration of telmisartan reduced systolic blood pressure and oxidative stress probably through the activation of PI3K/Akt/eNOS pathway and NO release in spontaneously hypertensive rats. *Physiol Res*. 2013;62:351–9.
- Mugabo P, Raji IA. Effects of aqueous leaf extract of *Asystasia gangetica* on the blood pressure and heart rate in male spontaneously hypertensive Wistar rats. *BMC Complement Altern Med*. 2013;13:283.
- Wu L, Juurlink BH. Increased methylglyoxal and oxidative stress in hypertensive rat vascular smooth muscle cells. *Hypertension*. 2002;39:809–14.
- Hultström M. Development of structural kidney damage in spontaneously hypertensive rats. *J Hypertens*. 2012;30:1087–91.

28. Pende A, Dallegri F. Renin-angiotensin antagonists: therapeutic effects beyond blood pressure control? *Curr Pharm Des.* 2012;18:1011–20.
29. Bakris G. Are there effects of renin-angiotensin system antagonists beyond blood pressure control? *Am J Cardiol.* 2010;105:21A–9.
30. Marchesi C, Paradis P, Schiffrin EL. Role of the renin-angiotensin system in vascular inflammation. *Trends Pharmacol Sci.* 2008;29:367–74.
31. Kobori H, Nangaku M, Navar LG, Nishiyama A. The intrarenal renin-angiotensin system: from physiology to the pathobiology of hypertension and kidney disease. *Pharmacol Rev.* 2007;59:251–87.
32. Unger T, Stoppelhaar M. Rationale for double renin-angiotensin-aldosterone system blockade. *Am J Cardiol.* 2007;100:25J–31.
33. Kim SM, Park S, Choue R. Effects of fermented milk peptides supplement on blood pressure and vascular function in spontaneously hypertensive rats. *Food Sci Biotechnol.* 2010;19:1409–13.
34. Sobajima M, Nozawa T, Ihori H, Shida T, Ohori T, Suzuki T, et al. Repeated sauna therapy improves myocardial perfusion in patients with chronically occluded coronary artery-related ischemia. *Int J Cardiol.* 2013;167:237–43.
35. Hwang S, Lee DH, Lee IK, Park YM, Jo I. Far-infrared radiation inhibits proliferation, migration, and angiogenesis of human umbilical vein endothelial cells by suppressing secretory clusterin levels. *Cancer Lett.* 2014;346:74–83.
36. Hsu YH, Chen YC, Chen TH, Sue YM, Cheng TH, Chen JR, et al. Far-infrared therapy induces the nuclear translocation of PLZF which inhibits VEGF-induced proliferation in human umbilical vein endothelial cells. *PLoS One.* 2012;7:e30674.
37. Akasaki Y, Miyata M, Eto H, Shirasawa T, Hamada N, Ikeda Y, et al. Repeated thermal therapy up-regulates endothelial nitric oxide synthase and augments angiogenesis in a mouse model of hindlimb ischemia. *Circ J.* 2006;70:463–70.
38. Meigs JB, Hu FB, Rifai N, Manson JE. Biomarkers of endothelial dysfunction and risk of type 2 diabetes mellitus. *JAMA.* 2004;291:1978–86.
39. Boulbou MS, Koukoulis GN, Makri ED, Petinaki EA, Gourgoulis KI, Germentis AE. Circulating adhesion molecules levels in type 2 diabetes mellitus and hypertension. *Int J Cardiol.* 2005;98:39–44.
40. Kressel G, Trunz B, Bub A, Hülsmann O, Wolters M, Lichtinghagen R, et al. Systemic and vascular markers of inflammation in relation to metabolic syndrome and insulin resistance in adults with elevated atherosclerosis risk. *Atherosclerosis.* 2009;202:263–71.
41. De Caterina R, Ghiadoni L, Taddei S, Virdis A, Almerigogna F, Basta G, et al. Soluble E-selectin in essential hypertension: a correlate of vascular structural changes. *Am J Hypertens.* 2001;14:259–66.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

